

| Red High priority | | |
|---|------------|-----------|
| Conditions | Short term | Long term |
| Alloimmune Thrombocytopenia (Foeto - Maternal/Neonatal) | | |
| Chronic inflammatory demyelinating polyradiculoneuropathy | | |
| Guillain Barre Syndrome | | |
| Haemolytic disease of the newborn | | |
| HSCT in primary immunodeficiencies | | |
| Immune thrombocytopenic purpura (acute and persistent, excluding chronic) | | |
| Kawasaki disease | | |
| Paraprotein - associated demyelinating neuropathy (IgM, IgG or IgA) | | |
| Primary immunodeficiencies | | |
| Specific antibody deficiency | | |
| Thymoma with immunodeficiency | | |
| Toxic epidermal necrolysis, Stevens Johnson syndrome | | |
| *Updated May 2018 | | |

| Blue Medium priority | | |
|---|------------|-----------|
| Conditions | Short term | Long term |
| Acquired red cell aplasia | | |
| Autoimmune congenital heart block | | |
| Autoimmune haemolytic anaemia | | |
| Autoimmune uveitis | | |
| Coagulation factor inhibitors (alloantibodies and autoantibodies) | | |
| Haemophagocytic syndrome | | |
| Immunobullous disease | | |
| Inflammatory myopathies | | |
| Multifocal motor neuropathy | | |
| Myasthenia gravis (including Lambert - Eaton myasthenic syndrome) | | |
| Necrotising (PVL - associated) staphylococcal sepsis | | |
| Post - transfusion purpura | | |
| Rasmussen syndrome | | |
| Secondary antibody deficiency (any cause) | | |
| Severe or recurrent clostridium difficile colitis | | |
| Staphylococcal Streptococcal toxic shock syndrome | | |
| Stiff person syndrome | | |
| Transplantation (solid organ) | | |

| Grey Low Priority | |
|--|--|
| Conditions | |
| Immune-mediated disorders with limited evidence of immunoglobulin efficacy | Presumed immune-mediated disorders with little or no evidence of efficacy |
| Acute disseminated encephalomyelitis (If high dose steroids have failed) | Acquired red cell aplasia NOT due to parvovirus B19 |
| Autoimmune encephalitis (including NMDA and VGKC antibodies, among others) | Acute idiopathic dysautonomia |
| Catastrophic antiphospholipid syndrome | Aplastic anaemia/pancytopenia |
| Cerebral infarction with antiphospholipid antibodies | Atopic dermatitis/eczema |
| Chronic ITP | Autoimmune neutropenia |
| CNS Vasculitis | Chronic facial pain |
| Complex regional pain syndrome | Diabetic proximal neuropathy |
| Intractable childhood epilepsy | Haemolytic uraemic syndrome |
| Neuromyotonia | PANDAS |
| Opsoclonus Myoclonus | Paraneoplastic disorders that are known not to be B- or T-cell mediated |
| Post exposure prophylaxis for viral or pathogenic infection if intramuscular injections is contraindicated, or treatment when hyper-immune Immunoglobulins are unavailable | POEMS |
| Pyoderma gangrenosum | SLE without secondary immunocytopenias (Including juvenile) |
| Systemic juvenile idiopathic arthritis | |
| Systemic vasculitides and ANCA disorders | |
| Urticaria (Severe, intractable) | |

| Black |
|--|
| Conditions |
| Immunodeficiency secondary to paediatric HIV infection |
| Autologous BMT |
| Adrenoleukodystrophy |
| Alzheimer's disease |
| Amyotrophic lateral sclerosis |
| Chronic fatigue syndrome |
| Critical illness neuropathy |
| Multiple sclerosis |
| Rheumatoid arthritis |
| Neonatal sepsis (prevention or treatment) |
| Sepsis in the intensive care unit not related to specific toxins or C. difficile |
| Asthma |
| Graves ophthalmopathy |
| IVF failure |
| Recurrent spontaneous pregnancy loss |